


## Control method: Aerial shooting of feral donkeys

### Assumptions:

- Best practice is followed in accordance with the standard operating procedure DON002.
- The shooter is competent and will make accurate decisions about whether the shot can be successfully placed. Competency also applies to the pilot who is required to provide the optimum target presentation for the shooter.
- Best practice is a shot to the head to render the animal immediately insensible followed by insurance shots to the chest.
- Group sizes are likely to be small (around 6 animals) although larger groups are sometimes found.

### PART A: assessment of overall welfare impact



<b>SCORE FOR PART A:</b>	<b>3-4</b>
Summary of evidence:	
Domain 1	No impact in this domain.
Domain 2	There is less opportunity to move away from the shooter compared with ground shooting. Exercise challenge is increased as there is likely to be a period of pursuit before being shot.
Domain 3	There is some potential for injury during helicopter pursuit. The wounding rate may be higher with aerial shooting (compared with ground shooting) because animals are shot whilst they are moving, however the range is likely to be much shorter and any wounded animals can be followed up quickly.
Domain 4	Aerial shooting forces donkeys to run and their flight response tends to be erratic resulting in the scattering of individuals. It therefore takes longer to locate and target individual animals because of this dispersal. The entire group is always killed therefore there are no long-term effects on social groups.
Domain 5	The presence of the helicopter will induce an escape response that includes running. This fear response is less than for horses, but still considered more than just a mild impact.

## PART B: assessment of mode of death – head shot

Time to insensibility (minus any lag time)				
Very rapid	Minutes	Hours	Days	Weeks
Level of suffering (after application of the method that causes death but before insensibility)				
No suffering	Mild suffering	Moderate suffering	Severe suffering	Extreme suffering

## PART B: assessment of mode of death – chest shot

Time to insensibility (minus any lag time)				
Very rapid	Minutes	Hours	Days	Weeks
Level of suffering (after application of the method that causes death but before insensibility)				
No suffering	Mild suffering	Moderate suffering	Severe suffering	Extreme suffering

<b>SCORE FOR PART B:</b>	<b>Head shot - A</b> <b>Chest shot - C</b>
Summary of evidence:	

Duration –	<p>With head shots, a properly placed shot will result in immediate insensibility<sup>1,2,3</sup>. A follow-up shot to ensure death ('insurance shot') is required in all cases.</p> <p>With chest shots, time to insensibility can range from seconds to a few minutes. The time to loss of consciousness and the time to death will depend on which tissues are damaged and, in particular, on the rate of blood loss and hence the rate of induction of cerebral hypoxaemia<sup>4</sup>. Loss of consciousness and death is likely to be quick when animals have been shot in the heart. 'Hydrostatic shock' (see below) may also contribute to rapid incapacitation and potentially rapid loss of consciousness with shots to the chest; however this effect seems to be variable and doesn't occur in all instances. 'Double tap' shots (two quick shots in succession) are always used with chest shots.</p>
Suffering –	<p>When animals are rendered insensible immediately with a well-placed head shot that causes adequate destruction of brain tissue there should be no suffering<sup>1</sup>.</p> <p>Animals that are chest shot and still conscious are likely to have a short period of suffering, though the extent of suffering will vary depending on which tissues are damaged and the rate of blood loss. During haemorrhage there is likely to be tachypnoea and hyperventilation, which, when severe, would indicate that there is a sense of breathlessness before the loss of consciousness<sup>4</sup>. Severe haemorrhage in humans is also associated with anxiety and confusion<sup>5</sup>.</p> <p>If chest shot animals are rendered insensible by the mechanism of 'hydrostatic shock' and they do not regain consciousness prior to death they are unlikely to suffer.</p>

## Summary

CONTROL METHOD:	<b>Aerial shooting of feral donkeys</b>
OVERALL HUMANENESS SCORE:	<p><b>Head shot – 3-4A</b></p> <p><b>Chest shot – 3-4C</b></p>
<p>Comments</p> <p><b>Wounding rates with aerial shooting</b></p> <p>Statistics on wounding rates for aerial culling of animals are not readily available. Information provided by Tim Fraser, Team Leader of the SA DEH aerial shooting team states that "Animals killed instantly by my team would be better than 90 % and wounded animals less than 5 %". He also explained that "In most cases an experienced shooter knows as he/she touches off the shot whether it is perfectly placed or not, and if there is any doubt, second or even third shots are on there way instantly".</p> <p>One published account of wounding rates during an aerial shooting cull of feral horses was found. This was in a report on the cull of feral horses in Guy Fawkes River National Park in 2000 prepared by English<sup>6</sup>. The cull occurred between 22 and 24 October 2000, during which time 606 horses were shot. One horse was found alive on 1st November despite having 2 bullet wounds in the killing zone. The report author states that 'many horses received 4 or more shots, but the great majority were killed by the first or second shot' (the actual numbers are not given in the report). Thirty-nine horses were examined after the cull on 2 and 10 November, and also 67 horses were examined by a veterinarian, and 'no evidence of indiscriminate killing away from the target zone was found'.</p> <p><b>Hydrostatic shock</b></p> <p>With shooting, in addition to the damage caused by the penetrating projectile, there is scientific evidence that organs can also be damaged by the pressure wave that occurs when a projectile enters</p>	

a viscous medium, a phenomenon known as 'hydrostatic shock'. Experimental studies on pigs and dogs demonstrate that a significant ballistic pressure wave reaches the brain of animals shot in an extremity such as the thigh<sup>8,9,10</sup>. It is hypothesised that damage to the brain occurs when the pressure wave reaches the brain from the thoracic cavity via major blood vessels but could also occur via acceleration of the head or by passage of the wave via a cranial mechanism<sup>11</sup>. It is also thought that hydrostatic shock may produce incapacitation more quickly than blood loss effects, however not all bullet impacts will produce a pressure wave strong enough to cause this rapid incapacitation<sup>12</sup>.

Anecdotal reports by hunters maintain that some species are more susceptible to this shock effect than others; however no studies were found that confirmed this. However there is some speculation that, if one of the mechanisms that contribute to the effect of hydrostatic shock and subsequent damage to the brain is caused by acceleration of the head, it is possible that some animals may be more resistant to the incapacitating effects of shooting. It is recognised that animals such as head-butting ruminants appear to be more resistant to concussion than humans and are thought to have a higher acceleration threshold which could make them more resistant to traumatic brain injury not only from externally imposed forces, accelerations and blunt force trauma but also from an internal ballistic pressure wave generated by a projectile<sup>13,14</sup>.

## Bibliography

1. American Veterinary Medical Association (2001). 2000 Report of the AVMA Panel on Euthanasia. *Journal of the American Veterinary Medical Association* **218**, 669-696
2. Gregory, N. (2004). *Physiology and behaviour of animal suffering*. (Blackwell: Oxford, UK).
3. Longair, J. et al. (1991). Guidelines for euthanasia of domestic animals by firearms. *Canadian Veterinary Journal* **32**, 724-726
4. Gregory, N.G. (2005). Bowhunting deer. *Animal Welfare* **14**, 111-116
5. Zajtchuk, R. (1995). Anesthesia and Perioperative Care of the Combat Casualty. Chapter 4 - Hemorrhage, Shock and Fluid Resuscitation. (Office of The Surgeon General at TMM Publications, Borden Institute, Walter Reed Army Medical Center: Washington, DC).at <[http://www.bordeninstitute.army.mil/published\\_volumes/anesthesia/ANfm.pdf](http://www.bordeninstitute.army.mil/published_volumes/anesthesia/ANfm.pdf)>
6. English, A.W. (2000). *Report on the cull of feral horses in Guy Fawkes River National Park in October 2000*. (Veterinary Clinical Sciences, Faculty of Veterinary Science, University of Sydney: Sydney).
7. Courtney, M. & Courtney, A. (2008). Scientific Evidence for Hydrostatic Shock. 0803.3051 at <<http://arxiv.org/abs/0803.3051>>
8. Suneson, A., Hansson, H. & Seeman, T. (1990). Pressure Wave Injuries to the Nervous System Caused by High-energy Missile Extremity Impact: Part I. Local and Distant Effects on the Peripheral Nervous System-A Light and Electron Microscopic Study on Pigs. *The Journal of Trauma* **30**,
9. Suneson, A., Hansson, H. & Seeman, T. (1990). Pressure Wave Injuries to the Nervous System Caused by High-energy Missile Extremity Impact: Part II. Distant Effects on the Central Nervous System-A Light and Electron Microscopic Study on Pigs. *The Journal of Trauma* **30**,
10. Wang, Q., Wang, Z., Zhu, P. & Jiang, J. (2004). Alterations of Myelin Basic Protein and Ultrastructure in the Limbic System at the Early Stage of Trauma-Related Stress Disorder in Dogs. *The Journal of Trauma* **56**,
11. Courtney, A. & Courtney, M. (2009). A thoracic mechanism of mild traumatic brain injury due to blast pressure waves. *Medical Hypotheses* **72**, 76-83
12. Courtney, A. & Courtney, M. (2007). Links between traumatic brain injury and ballistic pressure waves originating in the thoracic cavity and extremities. *Brain Injury* **21**, 657-662
13. Courtney, M. & Courtney, A. (2007). Sheep Collisions: the Good, the Bad, and the TBI. 0711.3804 at <<http://arxiv.org/abs/0711.3804>>
14. Shaw, N.A. (2002). The neurophysiology of concussion. *Progress in Neurobiology* **67**, 281-344