

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

BACKGROUND

Feral goats (*Capra hircus*) can have a significant impact on the environment and agricultural production and are a potential reservoir and vector of endemic and exotic diseases. Although often considered pests, feral goats are also an important resource, harvested commercially primarily for meat. Control methods include trapping, mustering, exclusion fencing, ground shooting and shooting from helicopters. Radio-collared 'Judas' goats are sometimes used during mustering to locate groups of feral goats. Refer to [GOA005 Use of Judas goats](#).

Mustering of feral goats is usually carried out on motor bike or horse with the aid of dogs, although helicopters or light aircraft can be used. Mustering is only effective and economic when goat densities are high. Shooting from a helicopter is a more effective method of quickly reducing feral goat populations. Refer to GOA002 Aerial shooting of feral goats.

Once mustered, the goats are usually sold to abattoirs for slaughter or, less commonly, for domestication, which offsets the costs of capture and handling. Where there is no market for them or where removal can be costly or impractical (eg in conservation areas or remote areas without access to transportation), the goats are sometimes destroyed by shooting in a holding yard or while being held in a mob by dogs.

This standard operating procedure (SOP) is a guide only. It does not replace or override the legislation that applies in the relevant state or territory jurisdiction. The SOP should only be used subject to the applicable legal requirements (including OH&S) operating in the relevant jurisdiction.

APPLICATION

- Mustering should only be used in a strategic manner as part of a coordinated program designed to achieve sustained effective control.
- Mustering is only efficient and economic when goat densities are high. Many landholders therefore opportunistically muster when they notice large groups of goats on their land. Commercial mustering can reduce goat densities to about one goat per square kilometre, after which the return for effort becomes uneconomic. However, with this approach, it is important that goats are not left to build up to undesirably high levels. Monitoring of goat numbers and resource degradation should determine the most suitable time to conduct goat control. In tablelands there might be times (usually in winter) when goats form larger mobs and so are more accessible.
- Mustering is relatively labour intensive compared to trapping and can be more stressful to the goats. In relatively flat and accessible country, mustering is usually done on horseback or motorbikes and with the aid of dogs. In rough, hilly country and more extensive areas, helicopters or light aircraft can be used to drive the goats towards a set of yards where a ground team completes the muster.
- Not all goats from an area will be mustered. Some animals might be left behind, including does with young kids, and others might become scattered.
- The use of radio-collared Judas goats to locate feral herds increases the effectiveness of mustering control operations, particularly when eradication is the aim of the program (refer to GOA005 Use of Judas goats).
- Trained herding/sheep dogs are sometimes used to muster feral goats.

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

- It is unacceptable to set a dog onto a goat with the intention of bringing it down, holding or attacking.
- To ensure that mustering, capture and handling is done with the least stress to the goats, operators must have a good knowledge of goat behaviour, including their movement patterns. They should also be familiar with the terrain they are to cover, so that dangerous areas can be avoided.
- Aircraft operators must ensure that their flying operations comply with requirements of the Civil Aviation Safety Authority.
- Shooting of goats should only be done by skilled operators who have the necessary experience with firearms and who hold the appropriate licences and accreditation.
- Storage and transportation of firearms and ammunition must comply with relevant legislation requirements.

ANIMAL WELFARE CONSIDERATIONS

Impact on target animals

- Mustering, capture and handling increase stress in feral goats, as they are not used to confinement or close contact with humans. As a result, these procedures can lead to mismothering, feeding disruption, social disruption, heat stress and also abortion in heavily pregnant females. Metabolic, nutritional and parasitic diseases and changes in environmental conditions are common causes of mortality and morbidity in confined feral goats.
- To avoid heat stress, mustering should be carried out in the cooler months.
- The tail end of the mob should set the pace rather than being forced to keep up with the leaders. Distances that the goats have to be mustered should be kept to a minimum (eg by using portable yards).
- Feral goats should be handled quietly and without force, to avoid panic and trampling.
- Goats that are severely injured during mustering or confinement must be killed quickly and humanely with a rifle shot to the head.
- Whenever possible, avoid mustering when females are kidding or have dependent young at foot. Unweaned kids might be left to die of starvation if their mothers are mustered and they are left behind.
- Although feral goats have been observed to breed at all times of the year, there are periods when the majority of kidding occurs (eg in southwest Queensland, kidding mainly occurs in May—June). Mustering should not be done in spring, as this is the time when there is a high proportion of does in late pregnancy or with kids at foot.
- Electric prods should not be used to assist in the handling of feral goats.
- Only trained working dogs are to be used to assist in the handling of feral goats. Trained sheep dogs such as kelpies are preferred, as they are not usually aggressive. As a precaution, muzzles can be fitted to dogs to prevent them causing bite injuries.
- Mixing unfamiliar groups or individual goats in yards can result in fighting, stress and injury. Normal social groups should be maintained whenever possible. There should be sufficient holding yards to avoid mixing different groups of stock.
- Only fit and healthy animals should be selected for transport. Heavily pregnant, very young or weak/sick/injured animals must either be destroyed, given proper veterinary assistance or transported at a later date when they are more suitable for transportation.

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

- The loading, transport, unloading, holding and slaughter of feral goats must be done with the minimum amount of stress, pain or suffering. Guidelines on these procedures can be found in relevant state or federal government guidelines. For example:
 - Pre-slaughter Management of Goats (Finn et al 2003)
 - Australian Standards and Guidelines for the Welfare of Animals — Land Transport of Livestock (AHA 2008).
- Also, the following Model Codes of Practice for the Welfare of Animals might be useful:
 - The Goat (SCAAHC 1991b)
 - Air Transport of Livestock (AAHQS 1986)
 - Rail Transport of Livestock (AAHQS 1983)
 - Sea Transport of Livestock (AAHQS 1987)
 - Livestock at Slaughtering Establishments (SCARM 2002b).
 - Guidelines on the export of live goats can be found in the Australian Standards for the Export of Livestock (DAFF 2011).

Impact on non-target animals

- Mustering is target specific and has minimal impact on other species.
- Dogs used for mustering must receive adequate care at all times. This includes food, water, shelter, safe and comfortable transportation, current vaccinations, worming, flea, tick and heartworm prevention, where appropriate. For more details refer to GEN002 The care and management of dogs used for pest animal control.

HEALTH AND SAFETY CONSIDERATIONS

- Care must be taken when handling goats, as they can carry diseases such as Q fever and scabby mouth (orf) that can affect humans and other animals. Routinely wash hands and other skin surfaces after handling goats or carcasses.
- Operators working with goats and goat carcasses are at risk of contracting Q fever. They can become infected when they inhale droplets of urine, milk, faeces or birth products from infected animals. Infection can also occur from inhalation of aerosols created during slaughter of infected animals, or dust from contaminated materials. Blood testing of personnel is recommended to assess previous exposure, followed by vaccination for susceptible individuals.
- The mustering, trapping and handling of feral goats is not without risk to the operators involved. A first aid kit should be carried at all times. Motor bike riders should wear helmets.
- Firearms are potentially hazardous. All people should stand well behind the shooter when goats are being shot. The line of fire must be chosen to prevent accidents or injury from stray bullets or ricochets.
- Firearm users must strictly observe all relevant safety guidelines relating to firearm ownership, possession and use. Firearms must be securely stored in a compartment that meets state legal requirements. Ammunition must be stored in a locked container separate from firearms.
- Adequate hearing protection should be worn by the shooter and others in the immediate vicinity. Repeated exposure to firearm noise can cause irreversible hearing damage.
- When shooting, safety glasses are recommended to protect eyes from gases, metal fragments and other particles.

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

EQUIPMENT REQUIRED

Yards

- Either portable or fixed holding yards can be used.
- The materials used must minimise the risks of injury or escape of goats once inside the enclosure. Projections such as loose wire or sharp edges likely to cause injury should be eliminated and fences should be secure and high enough to prevent goats escaping.
- Gates must be wide enough to allow the easy flow of animals, particularly bucks with large horns.
- Yards should be designed to minimise both dust and boggy conditions.
- If possible, yards should be positioned in a shady area with as much natural vegetation as possible.
- If mustering in extremes of climate (hot or cold) is unavoidable, shelter must be provided for goats. This is particularly important for young goats or animals in poor body condition during cold, windy and rainy conditions.
- Details of yard specifications and construction can be obtained from relevant state guidelines (eg Joshua 2003).

Light fixed wing aircraft or helicopter

- The aircraft must be suited to the purpose (eg Cessna 182, Hughes 500 helicopter, gyrocopter or ultralight fixed wing aircraft) and must be registered for the task of mustering.
- The pilot must be suitably licensed and hold the appropriate endorsements for aerial mustering of stock.

PROCEDURES

Mustering

- Goats should not be chased but moved steadily with the slowest animals setting the pace. Goats should never be driven to the point of collapse.
- Only muster the number of goats that can be comfortably handled. The fewer goats that are included in any one operation and the shorter the distance travelled, the less stress will be placed upon the animals.
- Heavily pregnant females, females with small kids and other goats, such as those in poor condition, should be allowed to drop out of musters if the safety and welfare of the animals is at risk. Also, if a female goat continually breaks away and will not move along with the group, it is possible that she has a dependent kid/s hidden somewhere. It is best to let her go and move on with the rest of the group.

Holding goats in yards

- Goats captured by mustering should be allowed at least 3–4 days rest with adequate shelter, food and water if they are to be transported for more than 8 hours. This will also allow them to become accustomed to lot-feeding before transport to a feedlot or depot. During this time they must be assessed daily for signs of injury, disease, loss of appetite, illness or distress. Account must be taken of their possible unwillingness to drink and eat from troughs.

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

- Goats should not be held in the holding yards for extended periods. If goats are being held for longer than four days, they should be drafted into a large holding paddock that contains adequate shelter, food and water.
- To minimise injury in yards, goats should be segregated into the following groups:
 - does with kids at foot
 - heavily pregnant does and small or young goats
 - bucks.

Loading and transporting goats

- Specific requirements for the land transport of goats can be found in Australian Standards and Guidelines for the Welfare of Animals — Land Transport of Livestock (AHA 2008).

Shooting of goats

- It might be necessary to humanely destroy goats by shooting in the following situations:
 - when there is no market for the captured goats (including smaller animals that are of no commercial value)
 - if goats have sustained serious injury during capture or in the holding yards
 - if there are dependent young that are separated from their mother
 - if there is a pre-existing disease or condition that would prevent the animal from being transported, slaughtered or domesticated.
- Shooting must be done to cause sudden and painless death with minimum distress to the animal. Only head shots are acceptable.
- The shooter should approach the animals in a calm and quiet manner. To prevent unnecessary agitation of the yarded goats, other people should keep away from the area until shooting is completed.
- To maximise the impact of the shot and to minimise the risk of misdirection, the range should be as short as possible.
- Never fire when the goat is moving its head. Be patient and wait until the goat is motionless before shooting. Accuracy is important to achieve a humane death. One shot should ensure instantaneous loss of consciousness and rapid death without resumption of consciousness.
- Shots must be aimed to destroy the major centres at the back of the brain near the spinal cord (see Diagram 1).
- This can be achieved by one of the following methods:

Head Shot

- The horn structures on adult goats make the temporal (side-on) or rear head shots the preferred points of aim. Shots to the front of the head can be used on kids, but this method is not recommended for mature goats as the brain is located well back in the skull.

Temporal position (side view)

- The firearm should be directed at the side of the head so that the bullet enters the skull midway between the eye and the base of the ear. The bullet should be directed horizontally.

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

Rear of the head

- The firearm should be aimed at the back of the head at a point between the base of the horns and directed towards the mouth.
- Death of shot animals should always be confirmed by observing the following:
 - absence of rhythmic respiratory movements
 - absence of eye protection reflex (corneal reflex) or 'blink'
 - a fixed, glazed expression in the eyes
 - loss of colour in mucous membranes (become mottled and pale without refill after pressure is applied).
 - If death cannot be verified, a second shot to the head should be taken immediately.
- When large numbers of animals are to be killed in the holding yard, provisions should be made to dispose of carcasses in an appropriate manner (ie by burying and/or burning). Numerous guidelines are available that describe disposal methods (eg Burton 1999, NSW EPA 2001, AHA 2007).

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

Recommended shot placements - Feral goat

Diagram 1




Diagram 2 - Side view (skeleton)




Diagram 3 - Head shot (rear)



MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

REFERENCES

1. AHA (Animal Health Australia) (2007). Operational Procedures Manual: Disposal (Version 3.0). Australian Veterinary Emergency Plan (AUSVETPLAN), Edition 3, Primary Industries Ministerial Council, Canberra, ACT.
2. AHA (2008). Australian Standards and Guidelines for the Welfare of Animals — Land Transport of Livestock. AHA, Canberra.
3. American Veterinary Medical Association (2007). AVMA Guidelines on Euthanasia (formerly The Report of the AVMA Panel on Euthanasia).
4. Department of Natural Resources and Mines (2001). Feral Goat (*Capra hircus*). NRM facts PA18. Land Protection, Department of Natural Resources and Mines, Queensland.
5. Biodiversity Group, Environment Australia (1999). Threat Abatement Plan for Competition and Land Degradation by Feral Goats. The Department of Environment and Heritage, through the National Threat Abatement Component of the Natural Heritage Trust.
6. AAHQS (Australian Agricultural Health and Quarantine Service) (1983). Model Code of Practice for the Welfare of Animals — Rail Transport of Livestock. Department of Primary Industry, Canberra.
7. AAHQS (1986). Model Code of Practice for the Welfare of Animals — Air Transport of Livestock. Department of Primary Industry, Canberra.
8. AAHQS (1987). Model Code of Practice for the Welfare of Animals — Sea Transport of Livestock. Department of Primary Industry, Canberra.
9. Burton R (1999). Humane Destruction and Disposal of Stock. Agnote DAI-136. NSW Agriculture, Orange.
10. Casburn G, Hacker R and Brill T (1999). Evaluation of Cooperative Feral Goat Harvesting/Control Techniques. Project 94/1. Vertebrate Pest Program. NSW Agriculture, Orange.
11. Australian Government Department of Agriculture, Fisheries and Forestry (DAFF) (2011). Australian Standards for the Export of Livestock. DAFF, Commonwealth of Australia.
12. English AW (2001). Code of Practice for the Capture and Transport of Feral Horses. A Report on the Management of Feral Horses in National Parks in New South Wales.
13. Finn J, Greenwood P and May T (2003). Pre-slaughter Management of Goats. Agfact A7.1.12. NSW Agriculture.
14. Henzell R (1984). Methods of Controlling Feral Goats. Factsheet. Agdex 573. Department of Agriculture, South Australia.
15. Jago B (1999). Feral Goat (*Capra hircus*) in Queensland. Pest Status Review Series – Land Protection. Department of Natural Resources and Mines, Queensland.
16. Joshua E (2003). Yard Design for Goats. Agfact A7.7.2. NSW Agriculture.
17. King D (1991). Goat Control Methods. Infonote no. 21/91. Agriculture Protection Board of Western Australia.
18. Longair JA, Finley GG, Laniel MA, MacKay C, Mould K, Olfert ED, Roswell H and Preston A (1991). Guidelines for euthanasia of domestic animals by firearms. Canadian Veterinary Journal 32:724-726.
19. O'Flynn M (1992). Animal Welfare Considerations. In: D Freudenberger (Ed), Proceedings of the National Workshop on Feral Goat Management: Planning for Action, Dubbo, NSW. Bureau of Resource Sciences, Canberra. Pp 39-55.
20. NSW EPA (Environment Protection Authority) (2001). Procedure for Dead Stock Disposal.

MUSTERING OF FERAL GOATS (GOA003) STANDARD OPERATING PROCEDURE

21. Parkes J, Henzell R and Pickles G (1996). Managing Vertebrate Pests: Feral Goats. Bureau of Resource Sciences and Australian Nature Conservation Agency, Australian Government Publishing Service, Canberra.
22. Primary Industries Ministerial Council (undated). Draft Model Code of Practice for the Welfare of Animals: Killing or Capture, Handling and Marketing of Feral Livestock Animals. CSIRO, Australia.
23. Ramsay BJ (1994). Commercial Use of Wild Animals in Australia. Bureau of Resource Sciences. Australian Government Publishing Service, Canberra.
24. SCAAHC (Standing Committee on Agriculture, Animal Health Committee) (1991a). Model Code of Practice for the Welfare of Animals: Feral Livestock animals — Destruction or Capture, Handling and Marketing. CSIRO Publishing, Australia.
25. SCAAHC (1991b). Model Code of Practice for the Welfare of Animals: The Goat. CSIRO Publishing, Australia.
26. SCARM (Standing Committee on Agriculture and Resource Management) (2002a). Model Code of Practice for the Welfare of Animals: Animals at Saleyards. SCARM Report 31. CSIRO Publishing, Collingwood.
27. SCARM (2002b). Model Code of Practice for the Welfare of Animals: Livestock at Slaughtering Establishments. SCARM Report 79. CSIRO Publishing, Collingwood.
28. Thompson J, Riethmuller J, Kelly D, Boyd-Law S and Miller E (1999). Technical Report on Feral Goat Management in South-west Queensland. Report to the Bureau of Rural Sciences. Queensland Department of Natural Resources.

The Centre for Invasive Species Solutions manages these documents on behalf of the Environment and Invasives Committee (EIC). The authors of these documents have taken care to validate the accuracy of the information at the time of writing. This information has been prepared with care but it is provided “as is”, without warranty of any kind, to the extent permitted by law.

If you have printed this document please ensure you regularly check <https://pestsmart.org.au> for the latest updates of these documents.